


Exposure	Personal Protective Equipment Used	Work Restrictions for Unvaccinated HCP
<p>Higher-risk: HCP who had prolonged<sup>1</sup> close contact<sup>2</sup> with a patient, visitor, or HCP with confirmed SARS-CoV-2 infection<sup>3</sup></p>	<ul style="list-style-type: none"> <li>• HCP not wearing a respirator or facemask<sup>4</sup></li> <li>• HCP not wearing eye protection if the person with SARS-CoV-2 infection was not wearing a cloth mask or facemask</li> <li>• HCP not wearing all recommended PPE (i.e., gown, gloves, eye protection, respirator) while performing an aerosol-generating procedure<sup>1</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Exclude from work for 14 days after last exposure.</li> <li>• Perform SARS-CoV-2 testing immediately (but not earlier than 2 days after the exposure) and, if negative, again 5-7 days after the exposure. Criteria for use of post-exposure prophylaxis are described <a href="#">elsewhere</a> .</li> <li>• Advise HCP to monitor themselves for fever or <a href="#">symptoms consistent with COVID-19</a>.</li> <li>• Any HCP who develop fever or <a href="#">symptoms consistent with COVID-19</a> should immediately contact their established point of contact (e.g., occupational health program) to arrange for medical evaluation and testing.</li> </ul>
<p>Lower-risk: HCP other than those with exposure risk described above</p>	<ul style="list-style-type: none"> <li>• N/A</li> </ul>	<ul style="list-style-type: none"> <li>• No work restrictions or testing.</li> <li>• Follow all <a href="#">recommended infection prevention and control practices</a>, including monitoring themselves for fever or <a href="#">symptoms consistent with COVID-19</a> and not reporting to work when ill. Any HCP who develop fever or <a href="#">symptoms consistent with COVID-19</a> should immediately self-isolate and contact their established point of contact (e.g., occupational health program) to arrange for medical evaluation and testing.</li> </ul>